

Dose-Dependent Decrease of Activation in Bilateral Amygdala and Insula by Lorazepam During Emotion Processing

Martin P. Paulus, MD; Justin S. Feinstein, BS; Gabriel Castillo, BS; Alan N. Simmons, PhD; Murray B. Stein, MD, MPH

Background: Functional neuroimaging may elucidate the pathophysiologic features of anxiety disorders and the site of action of anxiolytic drugs. A large body of evidence suggests that the amygdala and associated limbic structures play a critical role in the expression of anxiety and may be treatment targets for anxiolytic drugs.

Objective: To determine whether lorazepam dose-dependently attenuates blood oxygenation level-dependent functional magnetic resonance imaging (BOLD fMRI) activation in the amygdala and associated limbic structures during an emotion face assessment task.

Participants and Design: Fifteen healthy volunteers participated in a double-blind, placebo-controlled, randomized dose-response study. Subjects underwent imaging 3 times (at least a week apart) and were given either a single-dose placebo or 0.25 mg or 1.0 mg of lorazepam 1 hour prior to an MRI session. During fMRI, subjects completed an emotion face assessment task, which has been shown to elicit amygdala activation.

Main Outcome Measures: The BOLD-fMRI activation in amygdala, insula, and medial prefrontal cortex during the emotion face assessment task.

Results: Lorazepam significantly attenuated the BOLD-fMRI signal in a dose-dependent manner in bilateral amygdala and insula but not in the medial prefrontal cortex. Lorazepam did not affect the BOLD-fMRI signal in the primary visual cortex.

Conclusions: The current finding provides the first neuroimaging evidence of a dose-dependent change induced by an established therapeutic agent in brain regions known to be critical for the mediation of anxiety. This investigation may help to support the use of BOLD-fMRI with pharmacological probes to investigate the neural circuits underlying anxiety and the use of fMRI as a tool in the development of new anxiolytic agents.

Arch Gen Psychiatry. 2005;62:282-288

Author Affiliations: Laboratory of Biological Dynamics and Theoretical Medicine (Dr Paulus and Messrs Feinstein and Castillo) and Department of Psychiatry (Drs Paulus, Simmons, and Stein), University of California, San Diego; and Veterans Affairs San Diego Healthcare System (Drs Paulus and Stein).

ANXIETY DISORDERS ARE among the most common psychiatric disorders but comprise a heterogeneous group of conditions. Delineating the basic neurocircuitry underlying normal and pathologic anxiety¹ may help to better define anxiety disorders. The amygdala plays a critical role in normal fear conditioning and is implicated in the pathophysiologic mechanism of anxiety disorders.^{2,3} However, this structure is also important for other emotional information processing and behavior.⁴ Functional neuroimaging studies have shown amygdala activation in fear conditioning,⁵ reward-related processing,⁶ encoding of emotionally salient information,⁷ risk taking,⁸ processing positively valenced stimuli,⁹ and appetitive or aversive olfactory learning.¹⁰

In addition to the amygdala, a network of structures that includes the in-

sula, anterior cingulate gyrus, and medial prefrontal cortex is important to identify the emotional significance of a stimulus, generate an affective response, and regulate the affective state.¹¹ The insula has afferent and efferent connections to the medial and orbitofrontal cortex, anterior cingulate gyrus, and several nuclei of the amygdala.¹² Although insula activation has frequently been associated with disgust,¹³ there is increasing evidence of a broader role for this brain structure in emotion processing.¹⁴ Insula activation is thought to be involved in differential positive vs negative emotion processing,⁵ in particular fearful face processing,¹⁵ pain perception,^{16,17} and when individuals are asked to make judgments about emotions.¹⁸

Benzodiazepine derivatives are the most commonly prescribed antianxiety agents in clinical practice. Lorazepam, a 3-hydroxy-1,4-benzodiazepine, is rapidly and

readily absorbed, reaching peak concentrations in the blood proportional to the dose approximately 1 to 2 hours after oral administration.¹⁹ Previous neuroimaging investigation using 1 mg of intravenous lorazepam during a memory task found significant decreases in activation within the hippocampal, fusiform, and inferior prefrontal regions but no significant alterations in activation in the striate cortex.²⁰

The aim of this study was to use blood oxygenation level-dependent functional magnetic resonance imaging (BOLD fMRI) to test the hypothesis that lorazepam dose-dependently attenuates activation during an emotion face paradigm in the amygdala, insula, and medial prefrontal cortex. A modified emotion face assessment task was used to determine whether this attenuation would occur with negative emotion faces (angry and fear) but not with positive emotion faces (happy). Support for this hypothesis would provide a link between the clinical efficacy of lorazepam as an anxiolytic and the biological basis of γ -aminobutyric acid (GABA)-ergic modulation of the amygdala and limbic structures as key targets for anxiety circuitry.

Pharmacological fMRI is an emerging discipline with the potential to address a variety of neural systems and drug development questions.²¹ One important consideration with pharmacological fMRI is the differentiation between the drug's effect on neural tissue or on hemodynamic modulation. Some investigators have argued that pharmacological stimulation results in heterogeneous fMRI effects as opposed to global, homogeneous changes in BOLD signal. This argument is consistent with an action on specific neuronal receptor-based mechanisms.²² Moreover, focusing on a priori hypothesized areas may prevent false-positive findings.²³ Thus, the current study used primarily a region-of-interest approach.

METHODS

SUBJECTS

The University of California, San Diego, institutional review board approved the study procedures. All participants provided written informed consent and were paid for their participation. We studied 15 healthy, nonsmoking individuals (6 women and 9 men) aged 18 to 39 years (mean \pm SD, 27.6 \pm 1.4 years) with 12 to 18 years of education (mean \pm SD, 15.6 \pm 0.3 years). Participants did not have medical or psychiatric disorders as determined by medical history and diagnoses according to the Structured Clinical Interview for *Diagnostic and Statistical Manual of Mental Disorders, Revised Fourth Edition*. Each subject was required to consume less than 150 mg/d of caffeine and less than 14 caffeinated beverages per week. Subjects had no history of drug or alcohol abuse and did not report a prior use of benzodiazepines. All participants passed a urine drug screen. Subjects were instructed to maintain their regular bedtimes and wake times for 1 week before and throughout the study period. Individuals presented to the MRI facility between 8 AM and 4 PM with at least 1 week between studies.

STUDY DESIGN

The study was performed in a double-blind manner. Participants underwent each of 3 conditions in randomized order between 1 and 3 weeks apart. Lorazepam was chosen as the anxiolytic because of its well-documented behavioral effects and because it has a short metabolic half-life, which peaks at about 60 to 120 minutes after ingestion,²⁴ and no active metabolite, which eliminates the possible confounding effects of drug accumulation.²⁵ We selected doses that minimally disrupt performance of behavioral tasks and memory (≤ 1.0 mg) but are used to treat anxiety. Subjects arrived at the MRI facility 60 to 90 minutes prior to the MRI study. At arrival, subjects' vital signs were assessed and subjects orally received either placebo or 0.25 mg or 1.0 mg of lorazepam suspension mixed in diet decaffeinated cola.




Figure 1. Emotion face assessment task.²⁷ The target and 2 probe faces or ovals and circles were presented for 5 seconds. Blocks of each target emotion consisted of six 5-second angry, fearful, or happy face trials; blocks of ovals or circles consisted of two 5-second trials. Blocks of fixation crosses with varying time length were interspersed between the face and oval/circle blocks.

ASSESSMENTS AND fMRI PARADIGM

Subjects completed the Spielberger State-Trait Anxiety Inventory²⁶ (STAI) and the visual analog scales for anxiety, tension, alertness, and trembling before and after the MRI study. During fMRI, each subject was tested using a slightly modified version of the emotion face assessment task.²⁷ During each 5-second trial, a subject was presented with a target face (on the top of the computer screen) and 2 probe faces (on the bottom of the screen) and was instructed to match the probe with the same emotional expression to the target by pressing the left or right key on a button box (**Figure 1**). A block consisted of 6 consecutive trials in which the target face was either angry, fearful, or happy.²⁸ During the sensorimotor control task, subjects were presented with 5-second trials of ovals or circles in an analogous configuration and were instructed to match the shape of the probe to the target. Each block of faces and of the sensorimotor control task was presented 3 times in a pseudorandomized order. A fixation cross was interspersed between each block. For each trial, response accuracy and reaction time data were obtained.

IMAGE ACQUISITION

During the task, one BOLD-fMRI run was collected for each subject using a 1.5-T scanner (Siemens, Erlangen, Germany) (T2-weighted echoplanar imaging, time to repeat = 2000 milliseconds, echo time = 40 milliseconds, 64 \times 64 matrix, twenty 4-mm axial slices, 256 repetitions). During the same experimental session, a T1-weighted image (magnetization-




Figure 2. Behavioral effects of placebo and 0.25 mg and 1.0 mg of oral lorazepam. Lorazepam had no effect on response latency (A), the fraction of correct responses, or accuracy (B). Error bars indicate SE.

prepared rapid acquisition gradient echo, time to repeat = 11.4 milliseconds, echo time = 4.4 milliseconds, flip angle = 10°, field of view = 256 × 256, 1-mm³ voxels) was obtained for anatomical reference. For preprocessing, voxel time series were interpolated to correct for nonsimultaneous slice acquisition within each volume and were corrected for 3-dimensional motion.

IMAGE PROCESSING

All structural and functional image processing were done with the Analysis of Functional Neuroimages software package.²⁹ Echo-planar images were coregistered to the 128th image using a 3-dimensional coregistration algorithm. The time series of motion parameters was used to obtain a mean for these 6 parameters for each subject. Three motion parameters (d-roll, d-pitch, and d-yaw, indicating change in roll, pitch, and yaw directions) were used as nuisance regressors to account for motion artifacts. The 4 orthogonal regressors of interest were (1) happy, (2) angry, (3) fearful, and (4) circle/oval sensorimotor condition. These regressors were convolved with a modified gamma variate function to account for the delay and the dispersion brain response of the BOLD-fMRI signal due to hemodynamics.³⁰ Additional regressors were used to model residual motion in the roll, pitch, and yaw directions as well as baseline and linear trends. The Analysis of Functional Neuroimages program 3dDeconvolve²⁹ was used to calculate the estimated voxelwise response amplitude. A gaussian filter with 6 mm full width at half maximum was applied to the voxelwise percent signal change data to account for individual variations of the anatomical landmarks.

Data for each subject were normalized to Talairach coordinates. A priori regions of interest (defined by the Talairach Dae-

mon atlas³¹) in the bilateral amygdala, medial prefrontal cortex, primary visual cortex, and insula were used as masks. Based on these areas of interest, it was determined via simulations that a voxelwise a priori probability of .05 would result in a corrected clusterwise activation probability of .05 if a minimum volume of 128 μ L and 2 connected voxels (in the amygdala) or 512 μ L and 8 connected voxels (in all other regions of interest) was considered. The areas of interest were superimposed on each individual's voxelwise percent signal change brain image. Only activations within the areas of interest that also satisfied the volume and voxel connection criteria were extracted and used for further analysis. No clusters of voxels survived the cluster threshold parameters in the medial prefrontal cortex. Thus, planned comparisons using contrasts for repeated-measures designs were performed for only bilateral amygdala and insula when significant omnibus F values were found. The voxelwise percent signal change data were entered into a mixed-model analysis of variance with task contrast (face type vs circle/oval comparison condition) and dose (placebo or 0.25 or 1.0 mg of lorazepam) as fixed factors and subjects as a random factor. In addition, a whole-brain analysis was carried out to determine whether lorazepam affected nonhypothesized areas. Activations clusters were extracted and corrected for multiple comparisons using an a priori $P < .05$ threshold and a volume mask of greater than 1000 μ L.

STATISTICAL ANALYSIS

All behavioral analyses were carried out with SPSS version 10.0.³² A repeated-measures multivariate analysis of variance, with dose (placebo or 0.25 or 1.0 mg of lorazepam) as the within-subjects factor, was used to analyze the behavioral measures and neural activation patterns. Behavioral measures are reported as an interaction between dose and task type. Self-rating measures are reported as interactions between the drug and ratings before and after the MRI study. Correlational analyses were conducted for placebo administration by examining the relationship between self-rating scales or performance measures and activation in the bilateral insula and amygdala during viewing of angry, fearful, and happy faces.

RESULTS

BEHAVIORAL RESULTS

Subjects matched the probe face to the target face with nearly perfect accuracy (mean \pm SD, 97.0% \pm 0.7%), which was not affected by either the 0.25-mg or 1.0-mg dose of lorazepam ($F_{2,28} = 0.40$, $P = .60$, $\eta^2 = 0.03$). As shown in **Figure 2**, although latency was affected by face type (ie, longer for matching angry or fearful faces when compared with happy faces and circles or squares [$F_{3,12} = 48.8$, $P < .001$, $\eta^2 = 0.92$]), neither dose of lorazepam affected response latency ($F_{2,28} = 1.47$, $P = .25$, $\eta^2 = 0.09$).

As shown in **Figure 3**, lorazepam did not affect the level of anxiety, either measured by visual analog scales ($F_{2,22} = 1.71$, $P = .19$, $\eta^2 = 0.14$) or by the STAI ($F_{2,28} = 0.31$, $P = .73$, $\eta^2 = 0.02$). Moreover, visual analog ratings of tension ($F_{2,22} = 0.57$, $P = .57$, $\eta^2 = 0.05$) or trembling ($F_{2,22} = 1.82$, $P = .18$, $\eta^2 = 0.14$) did not significantly change across drug administration. However, individuals reported increased sleepiness after 1.0 mg of lorazepam, as evidenced by a dose by pre-post interaction ($F_{2,22} = 7.2$, $P = .004$, $\eta^2 = 0.40$) (Figure 2).




Figure 3. A, A high dose of lorazepam increased sleepiness (0=not at all; 10=extremely) at the 1.0-mg dose. B, Neither a low nor a high dose of lorazepam had a significant effect on state anxiety.

FUNCTIONAL NEUROIMAGING RESULTS

Irrespective of face type, subjects showed bilateral activation of the amygdala and anterior insula during the emotion face assessment task relative to the sensorimotor control task (see **Figure 4** and the **Table** for statistics). Lorazepam attenuated the activation in a dose-dependent fashion in the amygdala and insula (Table, **Figure 5A**). Specifically, activation in bilateral amygdala after administration of 1.0 mg of lorazepam was significantly lower than after 0.25 mg ($t_{14}=4.83$, $P<.001$) and significantly lower than after placebo administration ($t_{14}=2.85$, $P=.01$). There was no significant difference between placebo and 0.25 mg of lorazepam ($t_{14}=0.62$, $P=.54$). Similarly, activation in the bilateral insula after 1.0 mg of lorazepam was significantly lower than after 0.25 mg ($t_{14}=3.41$, $P=.004$) and significantly lower than after placebo administration ($t_{14}=3.63$, $P=.003$). There was no significant difference between placebo and 0.25 mg of lorazepam ($t_{14}=0.70$, $P=.50$). There was no significant effect of lorazepam on activation in the bilateral visual cortex for placebo or the 2 doses of lorazepam (Table, Figure 5B). A whole-brain analysis revealed significant effects of lorazepam in the right inferior temporal gyrus (coordinates: 43,-63,0), left insula (-27,7,-6), amygdala (-9,-11,-10), and fusiform gyrus (-29,-93,-13). In all areas, there was a significant attenuation of the signal in the 1.0-mg lorazepam condition relative to both placebo and 0.25 mg of lorazepam.




Figure 4. A, The blood oxygenation level–dependent functional magnetic resonance imaging effects (mean and SE) of placebo and 0.25 mg and 1.0 mg of oral lorazepam. B, Lorazepam dose-dependently attenuated the signal difference between the face-matching task and the sensorimotor control task in bilateral amygdala and insula. Numbers indicate Talairach z coordinates.

CORRELATIONS BETWEEN BEHAVIORAL AND FUNCTIONAL NEUROIMAGING RESULTS

There were no significant correlations between visual analog scales before administration or after imaging with the degree of activation in the bilateral insula or amygdala during task performance. Moreover, behavioral measures such as latency of response or accuracy of response during the emotion face assessment task did not correlate with the degree of activation in the bilateral insula or amygdala.

CONCLUSIONS

The results of this investigation show for the first time that a known anxiolytic, the benzodiazepine lorazepam, dose-dependently attenuates task-induced activation in bilateral amygdala and insula but has no effect

Table. Statistical Analyses of Regions of Interest

Effect	Amygdala			Insula			Visual Cortex		
	F Value	P Value	Partial Effect Size (η^2)	F Value	P Value	Partial Effect Size (η^2)	F Value	P Value	Partial Effect Size (η^2)
Laterality	1.65	.22	0.10	2.27	.15	0.14	0.25	.62	0.02
Dose	6.22	.006*	0.31	8.88	.001*	0.39	0.56	.58	0.04
Face†	1.39	.26	0.09	0.37	.69	0.03	2.62	.09	0.16
Side \times dose	0.57	.57	0.04	3.12	.06	0.18	0.26	.77	0.02
Side \times face	0.25	.78	0.02	0.10	.91	0.01	1.97	.16	0.12
Dose \times face	0.52	.72	0.04	0.68	.61	0.05	0.33	.86	0.02
Side \times dose \times face	1.18	.33	0.08	0.35	.84	0.02	1.00	.42	0.07

*Significant at $P < .05$.

†Target face: happy, fearful, or angry.




Figure 5. A, Percentage of signal activation during the face matching vs the sensorimotor comparison task at $z = -12$ during administration of placebo and 1.0 mg of oral lorazepam. Cross-hairs indicate the approximate center of mass of the amygdala as defined by the Talairach Daemon atlas.³¹ The bar indicates percent signal change from 0.2% to 1.0%. B, Blood oxygenation level-dependent functional magnetic resonance imaging percent signal difference of placebo and 0.25 mg and 1.0 mg of oral lorazepam in the visual cortex.

on the visual cortex. These effects were observed at doses of lorazepam that did not significantly affect behavior on this task and did not change levels of anxiety in these healthy (nonanxious) volunteers. In animals, a large body of evidence suggests that benzodiazepine agonists attenuate brain activity in the amygdala consistent with their

anxiolytic effects.³³ In humans, however, evidence of benzodiazepine effects on reducing amygdala activity had been missing.

A major focus in the field of anxiety research is the delineation of the basic neurocircuitry underlying normal and pathologic anxiety.¹ Amygdala activation occurs during fear conditioning, and altered amygdala activation has been implicated in the pathophysiologic mechanism of anxiety disorders.^{2,3} For example, individuals with social anxiety disorder³⁴ or posttraumatic stress disorder³⁵ show amygdala hyperresponsivity to fearful or angry faces. Moreover, patients with panic disorder have decreased benzodiazepine receptor binding in the left hippocampus and precuneus³⁶ as well as in the right orbitofrontal cortex and right insula.³⁷ The insular cortex appears to be important for subjective feeling states and interoceptive awareness.^{38,39} To our knowledge, this is the first study to show that an anxiolytic drug can dose-dependently reduce activation in the amygdala and insula, areas that are important for emotional processing in general and anxiety in particular. Thus, pharmacological fMRI with specific behavioral probes may prove especially useful for delineating how these brain structures are altered in anxiety disorders.

Lorazepam also attenuated activation in the fusiform gyrus. Although primarily recognized as a face-processing area,⁴⁰ this structure's activation patterns are related to the degree of conscious perception,⁴¹ anticipation,⁴² and identification of emotionally important visual clues.⁴³ Moreover, activation in the fusiform gyrus is altered in individuals with anxiety disorders,⁴⁴ and increased activation in the fusiform gyrus to fearful faces has been observed after cholinergic enhancement.⁴⁵ The amygdala can modulate early visual processing of emotional faces in the extrastriate cortex.¹⁵ Thus, lorazepam-induced attenuation of fusiform activity may be indirectly elicited by its down-regulation of the amygdala.

The behavioral effects of benzodiazepines in healthy, nonanxious volunteers have been mixed; for example, diazepam had no effect on fear-potentiated startle response,⁴⁶ whereas it impaired the recognition of anger and fear but not other emotional expressions.⁴⁷ Single administration vs repeated administration of lorazepam can induce loss of appetite, dizziness, and physical and mental

tiredness.^{48,49} Benzodiazepines dose-dependently disrupt learning and performance on acute vigilance tasks⁵⁰ and on a variety of memory tasks, presumably by reducing the normally facilitative effect of emotion on memory.^{51,52} Thus, behavioral effects may not provide robust indicators of potential anxiolytic efficacy in healthy nonanxious subjects, whereas changes in neural activation patterns induced by these substances may be informative. A low level of state anxiety may account for the lack of correlation between attenuation of the amygdala by lorazepam and changes in anxiety levels in our study. Future studies may include individuals with high trait anxiety (and ultimately anxiety disorders) to elucidate the relationship between anxiolytic effect and amygdala attenuation.

New drugs with novel anxiolytic mechanisms have been developed, including GABA A subreceptors,⁵³ metabotropic glutamate receptor agents,⁵⁴ and neuro-peptide modulators.⁵⁵ The current results provide a strong rationale for using BOLD fMRI imaging and amygdala- and/or insula-sensitive behavioral paradigms to determine whether these substances exert effects similar to those seen with standard anxiolytics. Furthermore, evaluation of the effects of other known classes of anxiolytics, such as long-term administration of selective serotonin reuptake inhibitors, should be undertaken to further validate this paradigm. The effect sizes (η^2) observed in this study provide proof in principle that small-scale (ie, sample sizes in the range of 15 to 30 subjects) BOLD fMRI studies can provide reasonably robust data to determine the site of action of putative anxiolytics. In summary, the current finding provides the first evidence of a dose-dependent change induced by an established therapeutic agent in brain regions known to be critical for the mediation of anxiety.

Submitted for Publication: March 29, 2004; final revision received July 27, 2004; accepted August 20, 2004.

Correspondence: Martin P. Paulus, MD, Department of Psychiatry, University of California, San Diego, 8950 La Jolla Village Dr, Suite C213, La Jolla, CA 92037-0985 (mpaulus@ucsd.edu).

Funding/Support: This work was supported by grants DA13186 and MH65413 from the National Institutes of Health, Bethesda, Md.

Acknowledgment: We acknowledge the invaluable help of Kelly Winternheimer, BS, and Thuy Le, BS.

REFERENCES

- Kent JM, Rauch SL. Neurocircuitry of anxiety disorders. *Curr Psychiatry Rep*. 2003;5:266-273.
- Rauch SL, Shin LM, Wright CI. Neuroimaging studies of amygdala function in anxiety disorders. *Ann N Y Acad Sci*. 2003;985:389-410.
- Charney DS. Neuroanatomical circuits modulating fear and anxiety behaviors. *Acta Psychiatr Scand Suppl*. 2003;38-50.
- LeDoux JE. Brain mechanisms of emotion and emotional learning. *Curr Opin Neurobiol*. 1992;2:191-197.
- Buchel C, Morris J, Dolan RJ, Friston KJ. Brain systems mediating aversive conditioning: an event-related fMRI study. *Neuron*. 1998;20:947-957.
- Breiter HC, Rosen BR. Functional magnetic resonance imaging of brain reward circuitry in the human. *Ann N Y Acad Sci*. 1999;877:523-547.
- Canli T, Zhao Z, Brewer J, Gabrieli JD, Cahill L. Event-related activation in the human amygdala associates with later memory for individual emotional experience. *J Neurosci*. 2000;20:RC99.
- Ernst M, Bolla K, Mouratidis M, Contoreggi C, Matochik JA, Kurian V, Cadet JL, Kimes AS, London ED. Decision-making in a risk-taking task: a PET study. *Neuropsychopharmacology*. 2002;26:682-691.
- Garavan H, Pendergrass JC, Ross TJ, Stein EA, Risinger RC. Amygdala response to both positively and negatively valenced stimuli. *Neuroreport*. 2001;12:2779-2783.
- Gottfried JA, O'Doherty J, Dolan RJ. Appetitive and aversive olfactory learning in humans studied using event-related functional magnetic resonance imaging. *J Neurosci*. 2002;22:10829-10837.
- Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception, I: the neural basis of normal emotion perception. *Biol Psychiatry*. 2003;54:504-514.
- Augustine JR. Circuitry and functional aspects of the insular lobe in primates including humans. *Brain Res Brain Res Rev*. 1996;22:229-244.
- Phillips ML, Young AW, Scott SK, Calder AJ, Andrew C, Giampietro V, Williams SC, Bullmore ET, Brammer M, Gray JA. Neural responses to facial and vocal expressions of fear and disgust. *Proc R Soc Lond B Biol Sci*. 1998;265:1809-1817.
- Phan KL, Wager T, Taylor SF, Liberzon I. Functional neuroanatomy of emotion: a meta-analysis of emotion activation studies in PET and fMRI. *Neuroimage*. 2002;16:331-348.
- Morris JS, Friston KJ, Buechel C, Frith CD, Young AW, Calder AJ, Dolan RJ. A neuromodulatory role for the human amygdala in processing emotional facial expressions. *Brain*. 1998;121:47-57.
- Gelnar PA, Krauss BR, Sheehee PR, Szeverenyi NM, Apkarian AV. A comparative fMRI study of cortical representations for thermal painful, vibrotactile, and motor performance tasks. *Neuroimage*. 1999;10:460-482.
- Peyron R, Laurent B, Garcia-Larrea L. Functional imaging of brain responses to pain: a review and meta-analysis (2000). *Neurophysiol Clin*. 2000;30:263-288.
- Gorno-Tempini ML, Pradelli S, Serafini M, Pagnoni G, Baraldi P, Porro C, Nicoletti R, Umiltà C, Nichelli P. Explicit and incidental facial expression processing: an fMRI study. *Neuroimage*. 2001;14:465-473.
- Kyriakopoulos AA, Greenblatt DJ, Shader RI. Clinical pharmacokinetics of lorazepam: a review. *J Clin Psychiatry*. 1978;39:16-23.
- Sperling R, Greve D, Dale A, Killiany R, Holmes J, Rosas HD, Cocchiarella A, Firth P, Rosen B, Lake S, Lange N, Routledge C, Albert M. Functional MRI detection of pharmacologically induced memory impairment. *Proc Natl Acad Sci U S A*. 2002;99:455-460.
- Salmeron BJ, Stein EA. Pharmacological applications of magnetic resonance imaging. *Psychopharmacol Bull*. 2002;36:102-129.
- Stein EA. fMRI: a new tool for the in vivo localization of drug actions in the brain. *J Anal Toxicol*. 2001;25:419-424.
- Stein EA, Pankiewicz J, Harsch HH, Cho JK, Fuller SA, Hoffmann RG, Hawkins M, Rao SM, Bandettini PA, Bloom AS. Nicotine-induced limbic cortical activation in the human brain: a functional MRI study. *Am J Psychiatry*. 1998;155:1009-1015.
- Busto UE, Kaplan HL, Wright CE, Gomez-Mancilla B, Zawertailo L, Greenblatt DJ, Sellers EM. A comparative pharmacokinetic and dynamic evaluation of alprazolam sustained-release, bromazepam, and lorazepam. *J Clin Psychopharmacol*. 2000;20:628-635.
- Rush CR, Higgins ST, Bickel WK, Hughes JR. Acute effects of triazolam and lorazepam on human learning, performance and subject ratings. *J Pharmacol Exp Ther*. 1993;264:1218-1226.
- Spielberger CD. *Manual for the State-Trait Anxiety Inventory (Form Y)*. Palo Alto, Calif: Consulting Psychologists Press; 1983.
- Hariri AR, Mattay VS, Tessitore A, Kolachana B, Fera F, Goldman D, Egan MF, Weinberger DR. Serotonin transporter genetic variation and the response of the human amygdala. *Science*. 2002;297:400-403.
- Ekman P, Levenson RW, Friesen WV. Autonomic nervous system activity distinguishes among emotions. *Science*. 1983;221:1208-1210.
- Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Comput Biomed Res*. 1996;29:162-173.
- Boynton GM, Engel SA, Glover GH, Heeger DJ. Linear systems analysis of functional magnetic resonance imaging in human V1. *J Neurosci*. 1996;16:4207-4221.
- Lancaster JL, Woldorff MG, Parsons LM, Liotti M, Freitas CS, Rainey L, Kochunov PV, Nickerson D, Mikiten SA, Fox PT. Automated Talairach atlas labels for functional brain mapping. *Hum Brain Mapp*. 2000;10:120-131.
- Norusis MJ. *SPSS Base System User's Guide*. Chicago, Ill: SPSS Inc; 1990.
- Menard J, Treit D. Effects of centrally administered anxiolytic compounds in animal models of anxiety. *Neurosci Biobehav Rev*. 1999;23:591-613.
- Stein MB, Goldin PR, Sareen J, Zorrilla LT, Brown GG. Increased amygdala activation to angry and contemptuous faces in generalized social phobia. *Arch Gen Psychiatry*. 2002;59:1027-1034.

35. Rauch SL, Whalen PJ, Shin LM, McNerney SC, Macklin ML, Lasko NB, Orr SP, Pitman RK. Exaggerated amygdala response to masked facial stimuli in post-traumatic stress disorder: a functional MRI study. *Biol Psychiatry*. 2000;47:769-776.
36. Bremner JD, Innis RB, White T, Fujita M, Silbersweig D, Goddard AW, Staib L, Stern E, Capiello A, Woods S, Baldwin R, Charney DS. SPECT [I-123]iomazenil measurement of the benzodiazepine receptor in panic disorder. *Biol Psychiatry*. 2000;47:96-106.
37. Malizia AL, Cunningham VJ, Bell CJ, Liddle PF, Jones T, Nutt DJ. Decreased brain GABA(A)-benzodiazepine receptor binding in panic disorder: preliminary results from a quantitative PET study. *Arch Gen Psychiatry*. 1998;55:715-720.
38. Craig AD. How do you feel? Interoception: the sense of the physiological condition of the body. *Nat Rev Neurosci*. 2002;3:655-666.
39. Critchley HD, Wiens S, Rotshtein P, Ohman A, Dolan RJ. Neural systems supporting interoceptive awareness. *Nat Neurosci*. 2004;7:189-195.
40. Kanwisher N, McDermott J, Chun MM. The fusiform face area: a module in human extrastriate cortex specialized for face perception. *J Neurosci*. 1997;17:4302-4311.
41. Andrews TJ, Schluppeck D, Homfray D, Matthews P, Blakemore C. Activity in the fusiform gyrus predicts conscious perception of Rubin's vase-face illusion. *Neuroimage*. 2002;17:890-901.
42. Chua P, Krams M, Toni I, Passingham R, Dolan R. A functional anatomy of anticipatory anxiety. *Neuroimage*. 1999;9:563-571.
43. Geday J, Gjedde A, Boldsen AS, Kupers R. Emotional valence modulates activity in the posterior fusiform gyrus and inferior medial prefrontal cortex in social perception. *Neuroimage*. 2003;18:675-684.
44. Bremner JD, Vythilingam M, Vermetten E, Southwick SM, McGlashan T, Staib LH, Soufer R, Charney DS. Neural correlates of declarative memory for emotionally valenced words in women with posttraumatic stress disorder related to early childhood sexual abuse. *Biol Psychiatry*. 2003;53:879-889.
45. Bentley P, Vuilleumier P, Thiel CM, Driver J, Dolan RJ. Cholinergic enhancement modulates neural correlates of selective attention and emotional processing. *Neuroimage*. 2003;20:58-70.
46. Baas JM, Grillon C, Bocker KB, Brack AA, Morgan CA III, Kenemans JL, Verbaten MN. Benzodiazepines have no effect on fear-potentiated startle in humans. *Psychopharmacology (Berl)*. 2002;161:233-247.
47. Zangara A, Blair RJ, Curran HV. A comparison of the effects of a beta-adrenergic blocker and a benzodiazepine upon the recognition of human facial expressions. *Psychopharmacology (Berl)*. 2002;163:36-41.
48. File SE, Bond AJ. Impaired performance and sedation after a single dose of lorazepam. *Psychopharmacology (Berl)*. 1979;66:309-313.
49. File SE, Bond AJ, Lister RG. Interaction between effects of caffeine and lorazepam in performance tests and self-ratings. *J Clin Psychopharmacol*. 1982;2:102-106.
50. Greenblatt DJ, Scavone JM, Harmatz JS, Engelhardt N, Shader RI. Cognitive effects of beta-adrenergic antagonists after single doses: pharmacokinetics and pharmacodynamics of propranolol, atenolol, lorazepam, and placebo. *Clin Pharmacol Ther*. 1993;53:577-584.
51. Buchanan TW, Karafin MS, Adolphs R. Selective effects of triazolam on memory for emotional, relative to neutral, stimuli: differential effects on gist versus detail. *Behav Neurosci*. 2003;117:517-525.
52. Blair RJ, Curran HV. Selective impairment in the recognition of anger induced by diazepam. *Psychopharmacology (Berl)*. 1999;147:335-338.
53. Whiting PJ. The GABAA receptor gene family: new opportunities for drug development. *Curr Opin Drug Discov Devel*. 2003;6:648-657.
54. Schoepp DD, Wright RA, Levine LR, Gaydos B, Potter WZ. LY354740, an mGlu2/3 receptor agonist as a novel approach to treat anxiety/stress. *Stress*. 2003;6:189-197.
55. Kent JM, Mathew SJ, Gorman JM. Molecular targets in the treatment of anxiety. *Biol Psychiatry*. 2002;52:1008-1030.